

## Multimodal Therapy (MMT) for Children with ADHD

### Program description:

These treatments target more than one setting with psychosocial interventions. For instance, many therapies intervene with both parents and teachers or children. In this analysis, all studies utilized either behavioral or cognitive-behavioral orientations.

Typical age of primary program participant: 8

Typical age of secondary program participant: N/A

### Meta-Analysis of Program Effects

Outcomes Measured	Primary or Secondary Participant	No. of Effect Sizes	Unadjusted Effect Sizes (Random Effects Model)			Adjusted Effect Sizes and Standard Errors Used in the Benefit-Cost Analysis					
			ES	SE	p-value	First time ES is estimated			Second time ES is estimated		
						ES	SE	Age	ES	SE	Age
Test scores	P	5	0.02	0.08	0.79	0.03	0.08	8	0.01	0.04	17
Disruptive behavior disorder symptoms	P	7	-0.29	0.13	0.02	-0.19	0.13	8	-0.08	0.05	13
Attention deficit hyperactivity disorder symptoms	P	9	-0.18	0.12	0.14	-0.08	0.12	8	-0.03	0.05	13

### Benefit-Cost Summary

The estimates shown are present value, life cycle benefits and costs. All dollars are expressed in the base year chosen for this analysis (2011). The economic discount rates and other relevant parameters are described in Technical Appendix 2.	Program Benefits				Costs	Summary Statistics				
	Partici-pants	Tax-payers	Other	Other Indirect		Total Benefits	Benefit to Cost Ratio	Return on Investment	Benefits Minus Costs	Probability of a positive net present value
	\$995	\$440	\$97	\$217	\$1,749	-\$8,343	\$0.21	n/e	-\$6,593	11%


### Detailed Monetary Benefit Estimates

Source of Benefits	Benefits to:					Total Benefits
	Partici-pants	Tax-payers	Other	In-direct		
Crime	\$0	\$9	\$26	\$4		\$39
Earnings via test scores	\$971	\$357	\$0	\$176		\$1,504
K-12 grade repetition	\$0	\$2	\$0	\$1		\$3
Health care costs for ADHD symptoms	\$11	\$34	\$34	\$17		\$97
Health care costs for disruptive behavior symptoms	\$13	\$38	\$37	\$19		\$107

### Detailed Cost Estimates

The figures shown are estimates of the costs to implement programs in Washington. The comparison group costs reflect either no treatment or treatment as usual, depending on how effect sizes were calculated in the meta-analysis. The uncertainty range is used in Monte Carlo risk analysis, described in Technical Appendix 2.	Program Costs			Comparison Costs			Summary Statistics	
	Annual Cost	Program Duration	Year Dollars	Annual Cost	Program Duration	Year Dollars	Present Value of Net Program Costs (in 2011 dollars)	Uncertainty (+ or - %)
	\$9,120	1	2010	\$950	1	2010	\$8,354	20%

Source: Based on therapist time, as reported in the treatment studies, as well as training costs and a flat fee for materials (e.g., manuals). Hourly therapist cost was based on the latest actuarial estimates of reimbursement by modality in WA State (DSHS).



### Multiplicative Adjustments Applied to the Meta-Analysis

Type of Adjustment	Multiplier
1- Less well-implemented comparison group or observational study, with some covariates.	1.00
2- Well-implemented comparison group design, often with many statistical controls.	1.00
3- Well-done observational study with many statistical controls (e.g., IV, regression discontinuity).	1.00
4- Random assignment, with some RA implementation issues.	1.00
5- Well-done random assignment study.	1.00
Program developer = researcher	0.64
Unusual (not "real world") setting	1.00
Weak measurement used	0.5

Adjustment factors were generated by examining studies for the treatment of children or adolescents with externalizing problems. Meta-regressions were conducted to test for the impact of different methodological factors on unadjusted effect size. Because research design rating and unusual setting were not significant predictors of effect size, multipliers were set to 1.0. A dummy variable representing involvement of a program developer in the research study was a statistically significant predictor ( $B = -.189, p = .056$ ), indicating that such studies had significantly more negative (i.e., larger) effect sizes than studies in which the developer was not involved. This coefficient was used to determine the 0.64 multiplier. Finally, we coded as weak measures outcomes that were based solely on the report of individuals who were involved in the intervention (either delivered it, as in the case of teachers, or received it, such as parents in a parenting program). Due to concern that such measures might be biased in favor of the programs reviewed, we utilized the standard Institute multiplier (0.5).

### Additional Notes

Some studies included in this analysis compared the program (MMT) to control conditions that did not consist of an active treatment. Because policymakers in Washington are interested in the impact of this program above and beyond currently implemented treatments (i.e., treatment as usual), we reduced the effect size of studies utilizing a no treatment or waitlist control group in half to reflect a smaller impact that would be expected if these studies compared MMT to treatment as usual.

### Studies Used in the Meta-Analysis

Abikoff, H., Hechtman, L., Klein, R. G., Weiss, G., Fleiss, K., Etcovitch, J., . . . Pollack, S. (2004). Symptomatic improvement in children with ADHD treated with long-term methylphenidate and multimodal psychosocial treatment. *Journal of the American Academy of Child & Adolescent Psychiatry, 43*(7), 802-811.

Chacko, A., Wymbs, B. T., Wymbs, F. A., Pelham, W. E., Swanger-Gagne, M. S., Girio, E., . . . O'Connor, B. (2009). Enhancing traditional behavioral parent training for single mothers of children with ADHD. *Journal of Clinical Child and Adolescent Psychology, 38*(2), 206-218.

Hechtman, L., Abikoff, H., Klein, R. G., Weiss, G., Respitiz, C., Kouri, J., . . . Pollack, S. (2004). Academic achievement and emotional status of children with ADHD treated with long-term methylphenidate and multimodal psychosocial treatment. *Journal of the American Academy of Child & Adolescent Psychiatry, 43*(7), 812-819.

Hechtman, L., Etcovitch, J., Platt, R., Arnold, L. E., Abikoff, H. B., Newcorn, J. H., . . . Wigal, T. (2005). Does multimodal treatment of ADHD decrease

## Studies Used in the Meta-Analysis

- other diagnoses? *Clinical Neuroscience Research*, 5(5-6), 273-282.
- Horn, W. F., Ialongo, N. S., Pascoe, J. M., Greenberg, G., Packard, T., Lopez, M., . . . Puttler, L. (1991). Additive effects of psychostimulants, parent training, and self-control therapy with ADHD children. *Journal of the American Academy of Child & Adolescent Psychiatry*, 30(2), 233-240.
- Klein, R. G., & Abikoff, H. (1997). Behavior therapy and methylphenidate in the treatment of children with ADHD. *Journal of Attention Disorders*, 2(2), 89-114.
- MTA Cooperative Group. (1999). A 14-month randomized clinical trial of treatment strategies for attention-deficit hyperactivity disorder. *Archives of General Psychiatry*, 56(12), 1073-1086.
- Pfiffner, L. J., Yee Mikami, A., Huang-Pollock, C., Easterlin, B., Zalecki, C., & McBurnett, K. (2007). A randomized, controlled trial of integrated home-school behavioral treatment for ADHD, predominantly inattentive type. *Journal of the American Academy of Child & Adolescent Psychiatry*, 46(8), 1041-1050.
- van der Oord, S., Prins, P. J. M., Oosterlaan, J., & Emmelkamp, P. M. G. (2007). Does brief, clinically based, intensive multimodal behavior therapy enhance the effects of methylphenidate in children with ADHD? *European Child & Adolescent Psychiatry*, 16(1), 48-57.